Referentiewaarden :

Leeftijd	Metanefrine (pmol/L)	Normetanefrine (pmol/L)	3MT (pmol/L)
< 1 dag	< 395	< 2811	< 479
1- 15 dagen	< 395	< 2725	< 460
15 dagen - 1 maand	< 401	< 2632	< 443
1 - 3 maanden	< 406	< 2304	< 377
3 – 6 maanden	< 421	< 1922	< 299
6- 9 maanden	< 431	< 1643	< 245
9 maanden – 1 jaar	< 441	< 1436	< 203
1 – 2 jaar	< 477	< 1026	< 126
2 – 3 jaar	< 497	< 901	< 108
3 – 15 jaar	< 370	< 737	< 78
15 – 18 jaar		< 557	
18 – 25 jaar		< 577	
25 – 30 jaar		< 601	
30 – 35 jaar		< 634	
35 – 40 jaar		< 677	
40 – 45 jaar		< 734	
45 – 50 jaar		< 804	
50 – 55 jaar		< 889	
55 – 60 jaar		< 992	
Vanaf 60 jaar		< 1113	
Vanaf 15 jaar	< 446		< 60

Bron:
Peitzsch M et al. Age-specific pediatric reference intervals for plasma free normetanephrine, metanephrine, 3-methoxytyramine and 3-O-methyldopa: Particular importance for early infancy. Clin Chim Acta. 2019 Jul;494:100-105.

Eisenhofer G et al. Reference intervals for LC-MS/MS measurements of plasma free, urinary free and urinary acid-hydrolyzed deconjugated normetanephrine, metanephrine and methoxytyramine. Clin Chim Acta. 2019 Mar;490:46-54. "